

The cosmological constant filter without big bang singularity

Florian Bauer

High Energy Physics Group, Dept. ECM, and Institut de Ciències del Cosmos
Universitat de Barcelona, Av. Diagonal 647, E-08028 Barcelona, Catalonia, Spain
Email: fbauerphysik@eml.cc

Abstract

In the recently proposed cosmological constant (CC) filter mechanism based on modified gravity in the Palatini formalism, gravity in the radiation, matter and late-time de Sitter eras is insensitive to energy sources with the equation of state -1 . This implies that finite vacuum energy shifts from phase transitions are filtered out, too. In this work we investigate the CC filter model at very early times. We find that the initial big bang singularity is replaced by a cosmic bounce, where the matter energy density and the curvature are finite. In a certain case this finiteness can be observed already on the algebraic level.

1 Introduction

The core of the old CC problem [1, 2, 3] is the huge hierarchy between the observed value $\Lambda_0 \sim 10^{-47} \text{ GeV}^4$ of the CC in the Λ CDM model and the value Λ inferred from theory. While the latter is difficult to calculate, it is usually expected to be dominated by the ultraviolet sector of the theory implying an enormous magnitude $|\Lambda| \gg \Lambda_0$. For example, the most reliable contribution to Λ comes from the electro-weak phase transition giving rise to the relation $|\Lambda| \sim (10^2 \text{ GeV})^4 \sim 10^{55} \Lambda_0$ [4]. A common way to deal with this problem is adding a CC counterterm, which subtracts off the huge vacuum energy density and permits a reasonable low-energy universe. The price to pay is the enormous finetuning of the counterterm, i.e. the precise choice of tens of digits, which is an undesired property for any theory. It is important to note that *replacing* the CC by a dynamical dark energy source generally does not avoid the old CC problem [5]. However, it may help modelling the late-time accelerated expansion and other cosmological epochs [6, 7, 8, 9, 10, 11, 12, 13].

Instead of removing the large CC by a static counterterm, we consider a more dynamical approach to neutralise the effects of Λ . It is easy to see that in standard general relativity the large CC would dominate the evolution of the universe at early times. Hence, one has to introduce nontrivial changes in gravity [14, 4, 15, 16] or in the matter sector [17, 18, 19] in order to permit a low curvature cosmos. Further ways to address the old CC problem can be found in Refs. [20, 21, 22, 23, 24, 25, 26], where, for instance, quantum gravity effects or self-adjustment mechanisms are important. Finally, neutralising the effect of vacuum energy was investigated in Refs. [27, 28, 29, 30, 31, 32] from different perspectives.

In this work we continue exploring the CC filter model proposed in Ref. [33]. This approach to solve the old CC problem is based on $f(R, Q)$ modified gravity in the Palatini formalism [34, 35, 36, 37, 38], where the gravity action is formulated in terms of the Ricci

scalar R and the squared Ricci tensor Q [39, 40, 41, 42]. Palatini theories have a more involved algebraic structure than the corresponding theories in the metric formalism [43, 44, 45, 46, 47, 48, 49, 50, 51], which permits the construction of a filter for vacuum energy. Moreover, in our setup we find only second order field equations just as in general relativity, thus extra degrees of freedom are avoided.

So far, only the region where the matter energy density ρ is smaller in magnitude than the CC Λ has been studied, which includes the radiation, matter and late-time de Sitter epochs. We found already on the algebraic level that energy sources with the equation of state (EOS) -1 do not contribute to the curvature at leading order, which is just a manifestation of the filter effect. In the following we will extend the analysis by discussing the very early cosmic epoch where ρ may be of the order of Λ or larger. As the main result we find that the filter model avoids the occurrence of an initial curvature singularity, which is instead replaced by a cosmic bounce with finite matter energy density and finite curvature. For a certain case, this finiteness can be seen also on the algebraic level. Finally, it is worth mentioning that the absence of an initial singularity has been observed also for other Palatini models in Refs. [52, 53], and a connection to loop quantum cosmology was suggested [54].

The paper is organised as follows: in Sec. 2 we briefly review how to solve the gravity field equations in the Palatini formalism, and the procedure will be applied to the CC filter model in Sec. 3. We solve the main equations and discuss the cosmological solutions in Sec. 4 before we conclude in Sec. 5.

In this work the speed of light c and the Planck constant $\hbar$ are set to unity, the signature of the metric is $(-1, +1, +1, +1)$. The tensor indices $a, \dots, f, m, n$ run from 0 to 3, whereas $i, j = 1, 2, 3$ are related to spatial coordinates.

2 $f(R, Q)$ modified gravity in the Palatini formalism

The CC filter mechanism is based on $f(R, Q)$ modified gravity¹ in the Palatini formalism, where R is the Ricci scalar and Q the squared Ricci tensor. In terms of the space-time metric g_{ab} and its determinant $g = \det g_{ab}$ the complete action is given by

$$\mathcal{S} = \int d^4x \left[\sqrt{|g|} \frac{1}{2} f(R, Q) \right] + \mathcal{S}_{\text{mat}}[g_{ab}, \phi] \quad (1)$$

with the matter fields ϕ in $\mathcal{S}_{\text{mat}}$ coupled minimally to g_{ab} . Furthermore, here the connection Γ_{bc}^a and the Ricci tensor R_{ab} are restricted to be symmetric,²

$$R_{ab} = \Gamma_{ab,e}^e - \Gamma_{eb,a}^e + \Gamma_{ab}^e \Gamma_{fe}^f - \Gamma_{af}^e \Gamma_{eb}^f \quad (2)$$

$$R = g^{ab} R_{ab}, \quad Q = R^{ab} R_{ab} = g^{ac} g^{bd} R_{ab} R_{cd}. \quad (3)$$

In the metric formalism the connection would be identified with the Levi-Civita connection

$$\Gamma_{bc}^a = \frac{1}{2} g^{ad} (g_{dc,b} + g_{bd,c} - g_{bc,d}), \quad (4)$$

and the Ricci tensor and all derived quantities would be functionals of the metric only. Differently, the Palatini formalism treats the connection as an independent object in the

¹The Palatini gravity model discussed in this work does not introduce extra degrees of freedom. Therefore, Ostrogradski-type [55] instabilities do not occur despite the squared Ricci tensor Q in the action.

²See Ref. [56] for generalisations.

beginning. Therefore, the variational principle $\delta\mathcal{S} = 0$ yields two equations of motion for gravity, one for the metric,

$$f_R R_m^n + 2f_Q R_m^a R_a^n - \frac{1}{2} \delta_m^n f = T_m^n, \quad (5)$$

and another one for the connection,

$$\nabla_a \left[\sqrt{|g|} (f_R g^{mn} + 2f_Q R^{mn}) \right] = 0, \quad (6)$$

where ∇_a denotes the covariant derivative in terms of the yet unknown connection Γ_{bc}^a and $f_{R,Q}$ are the partial derivatives of f with respect to R and Q , respectively.

In the following we briefly summarise how to solve these equations, and we point the reader to Refs. [33, 57] for a detailed exposition. Since we focus on the background cosmology, the matter sector can be described by a perfect fluid with the energy-stress tensor given by

$$T_m^n = (\rho + p) u_m u^n + (p - \Lambda) \delta_m^n, \quad (7)$$

where ρ is the energy density and p the pressure of ordinary matter with the 4-velocity vector u_m normalised by $u_m u^m = -1$. Λ denotes the large CC vacuum energy density.

From Eq. (5) two algebraic equations for the two unknowns R and Q can be derived, the first one is the trace of (5)

$$f_R R + 2Q f_Q - 2f = T, \quad (8)$$

where $T = T_m^m$. And the second equation follows from treating (5) as a matrix equation for the Ricci tensor $\hat{P} := R_m^n$ in matrix form,³

$$(2f_Q)^2 \left(\hat{P} + \frac{1}{4} \frac{f_R}{f_Q} \hat{I} \right)^2 = X^2 \hat{I} + 2f_Q (\rho + p) \widehat{u_m u^n} \quad (9)$$

with

$$X^2 := 2f_Q(p - \Lambda) + f_Q f + \frac{1}{4} f_R^2, \quad Y := X - c_b \cdot \sqrt{X^2 - 2f_Q(\rho + p)}, \quad (10)$$

The quadratic equation (9) has the solutions

$$c_a \cdot 2f_Q \left(\hat{P} + \frac{1}{4} \frac{f_R}{f_Q} \hat{I} \right) = X \hat{I} + Y \widehat{u_m u^n}, \quad (11)$$

where $c_{a,b} = \pm 1$ and the signs of the roots in X and Y follow from requiring that the final solution of the model solves Eq. (5). The trace of (11) yields

$$c_a (2f_Q R + 2f_R) = 4X - Y, \quad (12)$$

and at this point we simply eliminate all roots and ± 1 factors by squaring Eq. (12) twice,

$$\left((2f_Q R + 2f_R)^2 + 8X^2 + 2f_Q(\rho + p) \right)^2 = 36X^2 (2f_Q R + 2f_R)^2. \quad (13)$$

Using (8) and (13), the Ricci scalar R and the squared Ricci tensor Q can be determined as algebraic functions of ρ , p and Λ .

³Here, $\hat{I} = \delta_m^n$ is the identity matrix, and $\hat{P}^2 = R_m^a R_a^n$.

Next, we derive the connection Γ with the help of the auxiliary metric h_{mn} defined by

$$\sqrt{|h|}h^{mn} = \sqrt{|g|}g^{ma} \left(f_R \hat{I} + 2f_Q \hat{P} \right)_a^n, \quad (14)$$

which allows writing Eq. (6) in the form

$$\nabla_a[\Gamma] \left[\sqrt{|h|}h^{mn} \right] = 0. \quad (15)$$

Since $\hat{P}$ follows from Eq. (11), one finds after some algebra that h_{mn} is related to the “physical” metric g_{mn} by the disformal transformation [57]

$$h_{mn} = \Omega \left(g_{mn} - \frac{L_2}{L_1 - L_2} u_m u_n \right), \quad h^{mn} = \Omega^{-1} \left(g^{mn} + \frac{L_2}{L_1} u^m u^n \right), \quad (16)$$

where

$$\Omega := \sqrt{|L_1(L_1 - L_2)|}, \quad L_1 := c_a X + \frac{1}{2} f_R, \quad L_2 := c_a Y. \quad (17)$$

Finally, the Palatini connection Γ is just the solution of (15), i.e. the Levi-Civita connection of h_{mn} ,

$$\Gamma_{bc}^a[h] = \frac{1}{2} h^{ad} (h_{dc,b} + h_{bd,c} - h_{bc,d}) \quad (18)$$

with h_{mn} given in (16). This result solves Eq. (6) and eventually defines the Ricci tensor and all quantities derived from it. Note that h_{mn} does not contain space-time derivatives because it is a function of only g_{ab} , ρ , p and Λ . Hence, R_{ab} , R and the other quantities calculated according to Eqs. (2) and (3) contain derivatives of at most second order of the metric g_{ab} . Consequently, in this modified gravity theory in the Palatini formalism one has to solve the second order differential equation

$$R[\Gamma] = R(\rho, p, \Lambda), \quad (19)$$

where the right-hand side follows from the algebraic solution of Eqs. (8) and (13), whereas the left-hand side is given in Eq. (3) using Γ from Eq. (18). In the next section, we will apply this procedure to the CC filter model.

3 CC filter model

In Ref. [33] a modified gravity model was constructed that filters out vacuum energy contributions (i.e. everything with the EOS $p = -\rho$) to the CC. It is defined by the following action functional

$$f(R, Q) = \kappa R + z, \quad z := \beta \left(\frac{R^{\frac{2}{3}}}{B} \right)^m, \quad B := R^2 - Q, \quad (20)$$

with the dimensionful parameters κ and β and the positive number m . Note that in this setup, the term z is never considered to be a little correction to κR , instead both terms in f act in collaboration. From here on, we will use B instead of Q . With the partial derivatives of f , $f_R = (\kappa R + \frac{2}{3} m z)/R - 2f_Q R$, $f_Q = m z/B$, and the trace $T = -4\Lambda + 3p - \rho$ of the energy-stress tensor (7), Eq. (8) becomes

$$\gamma z = \kappa R - 4\Lambda + 3p - \rho, \quad \gamma := -2 - \frac{4}{3}m, \quad (21)$$

while Eq. (13) can be written as

$$0 = \kappa R + r - \frac{2}{9} m z \left(\frac{B}{R^2} \right) \left[1 + \frac{3}{m z} (3\kappa R + 2r) + \frac{9}{4(m z)^2} ((\kappa R)^2 - r^2) \right] \quad (22)$$

$$+ \frac{8}{27} m z \left(\frac{B}{R^2} \right)^2 \left(1 + \frac{3\kappa R}{2(m z)} \right)^2.$$

In the last equation we introduced the variable $r := \rho + p$, which vanishes identically for CC contributions.

For matter energy densities ρ and curvatures κR smaller in magnitude than the large CC, the term $\gamma z \approx -4\Lambda$ in (21) is approximately constant and Eq. (22) determines the curvature as a function of r alone. Therefore, contributions to the CC and shifts in the vacuum energy density are filtered out. This limit of constant $z = \mathcal{O}(\Lambda)$ covers most of the radiation epoch, the matter era and the late-time de Sitter phase, and it has been analysed analytically in Ref. [33] by neglecting strongly suppressed terms like $\mathcal{O}(r/z)$.

In the following we complement our previous results by investigating the very early universe, where r may be of the same order as Λ or above. Consequently, z in Eq. (21) is not constant anymore and all terms in Eq. (22) must be taken into account. It is convenient to work with dimensionless variables, where energy densities and the curvature term κR are normalised by the cosmological constant (4Λ),

$$y := \frac{\kappa R}{4\Lambda}, \quad x := \frac{r}{4\Lambda} = \frac{\rho + p}{4\Lambda}. \quad (23)$$

Moreover, let us define

$$m_z := \frac{m \cdot z}{4\Lambda} = \frac{m}{-\gamma} (1 - y - x \cdot \omega_m) \quad \text{with} \quad \omega_m := \frac{3\omega - 1}{\omega + 1}, \quad (24)$$

where $\omega = p/\rho$ denotes the matter EOS. Finally, we introduce the variable

$$W := \frac{R^2}{B} = y^{\frac{4}{3}} (1 - y - x \cdot \omega_m)^{\frac{1}{m}} \cdot \delta, \quad (25)$$

where B was eliminated on the right-hand side by using the definition of z in Eq. (20). The constant δ is completely fixed by the model parameters and the large vacuum energy density Λ ,

$$\delta := \left(\frac{2m}{-9\gamma} \right) \left(\frac{4\Lambda}{\rho_e} \right)^{\frac{7}{3}}, \quad \text{where} \quad (\rho_e)^{\frac{7}{3}} := \kappa^{\frac{4}{3}} \left(\frac{2}{9} m \frac{4\Lambda}{-\gamma} \right) \left(\frac{-\gamma\beta}{4\Lambda} \right)^{\frac{1}{m}}. \quad (26)$$

According to our earlier results in Ref. [33], δ is very large in magnitude because ρ_e is of the order of the late-time critical energy density in the limit $\gamma z \approx -4\Lambda$. Expressed in terms of the dimensionless variables the main equation (22) reads

$$0 = y + x - \frac{2}{9} \frac{m_z}{W} \left[1 + \frac{3(3y + 2x)}{m_z} + \frac{9(y^2 - x^2)}{4m_z^2} \right] + \frac{8}{27} \frac{m_z}{W^2} \left[1 + \frac{3y}{2m_z} \right]^2. \quad (27)$$

For the late-time solutions mentioned above we find $m_z = -m/\gamma = \mathcal{O}(1)$, $|y|, |x| \ll 1$ and $|W| \gg 1$. Therefore, Eq. (27) simplifies considerably,

$$y + x - \frac{2}{9} \frac{m_z}{W} = 0, \quad (28)$$

indicating $\kappa R \rightarrow \rho_e$ for vanishing matter $r \rightarrow 0$, which justifies introducing ρ_e in Eq. (26).

In the next section, we solve the complete equation (27) requiring a numerical treatment. In practise, we work along the following procedure. First, we consider a spatially flat cosmological background with the expansion described by the scale factor $a(t)$ as a function of cosmological time. In Cartesian coordinates ($i = 1, 2, 3$) the non-zero components of the metric are given by $g_{00} = -1$, $g_{ii} = a^2(t)$, and in the stress tensor we use $u_m = \delta_m^0$. Hence, the auxiliary metric in (16) has only the diagonal components

$$h_{00} = \Omega \left(\frac{-L_1}{L_{12}} \right), \quad h_{ii} = \Omega a^2(t), \quad L_{12} := L_1 - L_2, \quad (29)$$

which determine via (18) the symmetric components of the Palatini connection

$$\begin{aligned} \Gamma_{00}^0 &= \frac{\dot{x}}{4} \left(3 \frac{L_1'(x)}{L_1} - \frac{L_{12}'(x)}{L_{12}} \right) \\ \Gamma_{ii}^0 &= \frac{a^2}{L_1} \left(\frac{\dot{a}}{a} L_{12} + \frac{\dot{x}}{4} \left[L_{12}'(x) + \frac{L_{12} L_1'(x)}{L_1} \right] \right) \\ \Gamma_{i0}^i &= \frac{\dot{a}}{a} + \frac{\dot{x}}{4} \left(\frac{L_{12}'(x)}{L_{12}} + \frac{L_1'(x)}{L_1} \right). \end{aligned} \quad (30)$$

Here, the overdot indicates a derivative with respect to the time coordinate t , whereas the prime denotes the partial derivative with respect to x . For simplicity we consider matter with a constant EOS ω , which obeys the standard conservation equation $\dot{\rho} + 3\frac{\dot{a}}{a}\rho(1+\omega) = 0$ implying $\rho \propto a^{-3(1+\omega)}$ and $\dot{x} = -3\frac{\dot{a}}{a}(1+\omega)x$ via $x = (1+\omega)\rho/(4\Lambda)$. Consequently, the Ricci scalar $R(t)$ calculated via (3) from the connection (30) is a function of a , $\dot{a}$ and $\ddot{a}$, whereas the algebraic solution $y(x) = \kappa R/(4\Lambda)$ of Eq. (27) is only a function of a . For obtaining $a(t)$ we solve numerically the differential equation (19),

$$\frac{R(t)}{H_0^2} = K \cdot y(x(t)) \quad \text{with} \quad K := \frac{4\Lambda}{\kappa H_0^2}, \quad (31)$$

where $H_0 = \frac{\dot{a}}{a}(t_0)$ is the Hubble rate at the arbitrary initial time t_0 . In Ref. [33] we found $(-\kappa)$ to be of the order of the inverse Newton's constant G indicating $\kappa H_0^2 \sim \rho_e$ by identifying H_0 with the present Hubble rate and using the late-time solution (28). In this case, K is roughly the square root of δ in (26), its exact value is fixed when we specify H_0 and $x(t_0)$ at the initial⁴ time t_0 . Numerically, we first solve the equation⁵ $R^2/B = W(y(x))$ for $\ddot{a}(t_0)$, where R^2/B has to be calculated from (30) at $t = t_0$ and $x = x(a_0)$. Then, with $\ddot{a}(t_0)$ the values of $R(t_0)$ and $y(x(t_0))$ in Eq. (31) are known, which yields the numerical value of K and Eq. (31) can be solved to determine the evolution of the scale factor $a(t)$ for all times t .

4 Solutions

Here, we study the cosmological evolution resulting from the differential equation (31). Thereby we use the solutions to the main equation (27), which relate the curvature y

⁴Alternatively, one could start with given values of κ and $x(t_0)$ and infer H_0 from it. This is analogous to finding H_0 from the Friedmann equation $3H_0^2 = 8\pi G \cdot \rho(t_0)$ in general relativity.

⁵We use this equation in terms of arbitrary t to determine the correct choice of $c_{a,b}$ in (10) and (17) after Eq. (31) has been solved for $a(t)$.




Figure 1: The two branches of the function $y(x)$ from Eq. (27) relating the curvature term $y = \kappa R/(4\Lambda)$ to the matter variable $x = r/(4\Lambda)$ for the case $\delta > 0$, $\Lambda < 0$, EOS $\omega = \frac{1}{3}$.

to the matter energy density and pressure in x as given in (23). We do not restrict the magnitudes of x and y , however, in order to be roughly compatible with our existent universe, we consider only solutions $y(x)$ with both a final accelerating phase, $x \rightarrow 0$, and a high energy decelerating epoch, $y \approx -x$. Moreover, for the numerical examples below we set $m = 1/3$ in Eq. (20), because this parameter has only little influence on the qualitative behaviour of the solutions. With these preliminaries, there are four qualitatively different cases to study corresponding to $\Lambda \leq 0$ and $\delta \leq 0$. Note that the sign of δ can be adjusted by a suitable choice of the parameter β in the action functional (20).

1. The absence of an initial singularity can be seen best in the case $\delta > 0$ and $\Lambda < 0$ implying $\rho_e < 0$ via Eq. (26) and $\gamma < 0$ from (21). In this case the matter variable $x = r/(4\Lambda)$ is negative because of $r = \rho + p \geq 0$. Numerically, we do not find solutions for $y < 0$, that satisfy our requirements given above. This can be inferred also from Eq. (28), where all terms would be negative. On the other hand, for $y > 0$ (or $\kappa R < 0$) we find Eq. (28) as a good low-energy description with $y(x \rightarrow 0) \rightarrow \rho_e/(4\Lambda)$ and $y \approx -x$ for $|\rho_e| \ll |x| \ll 1$. However, before $y \rightarrow 1$, the matter variable $|x|$ becomes smaller again and eventually vanishes for finite $y < 1$ as illustrated in Fig. 1. As a result, $|x| \propto a^{-3(\omega+1)}$ and y are bounded from above signalling the absence of the initial singularity, where x or y would diverge. Analytically, this boundedness can be understood from Eqs. (24) and (25) indicating that for sufficiently large values of y or $|x|$ the variables W and m_z vanish. Since they appear in the denominators in our main equation (27), it is clear that neither $|x|$ nor y can become larger than 1. After solving the differential equation (31) for the scale factor $a(t)$, we observe that not the whole phase-space curve $y(x)$ in Fig. 1 is realised, but only a smaller part of it. This behaviour can be explained by observing that the function $L_1 - L_2$ vanishes at $x \approx -0.043$ as shown in Fig. 2. Since $L_1 - L_2$ appears in the denominator of the connection components (30) and therefore in the Ricci scalar $R(t)$, the equation (31)




Figure 2: The functions $L_1(x)$ and $L_{12}(x)$ in the connection components in Eqs. (30) for the case $\delta = 5 \cdot 10^6$, $\Lambda < 0$ and two values of the EOS $\omega = \frac{1}{3}, 0$. Notice the zero of L_{12} at $x \approx -0,043$, where the cosmic bounce occurs.

can be fulfilled only if $\dot{a}/a \propto \dot{x}/x$ vanishes at the zero of $L_1 - L_2$ because $y(x(t))$ on the right-hand side of (31) is always finite. Accordingly, $\dot{a} = 0$ indicates the occurrence of a cosmic bounce, which is shown in the plots for the scale factor $a(t)$, the Hubble rate $H = \dot{a}/a$ and the deceleration $q(t)$ in Fig. 3. Also, we can confirm the analytical results from Ref. [33], where it was found that at early times the universe behaves like a radiation- or dust-dominated cosmos depending on the matter EOS ω . Thus for $\omega = \frac{1}{3}$ or 0 the deceleration q approaches the value 1 or $\frac{1}{2}$, respectively. In this era, $y \approx -x$ is a good approximation. Finally, at late times ($x \rightarrow 0$) we find a de Sitter cosmos with $y \rightarrow \rho_e/(4\Lambda)$.

2. The second case we consider is $\delta < 0$ and $\Lambda > 0$ implying $x > 0$ and $\rho_e < 0$. Here, the only reasonable solution requires $y, \kappa R < 0$. In contrast to the first case, the variables x and $|y|$ are not bounded from above because W and m_z in Eqs. (24) and (25) do not vanish. Instead we find $y \rightarrow -x$ for large $x \rightarrow \infty$, see Fig. 4. Also, the functions L_1 and $L_1 - L_2$ do not vanish in this region. Nevertheless, the cosmic evolution inferred from the numerical solution of (31) shows a cosmic bounce, where the scale factor $a(t)$ and the matter and curvature variables $x, |y|$ remain finite, see Fig. 5. As before, the universe shows a radiation- or matter-dominated expansion behaviour with $y \approx -x$ and a final de Sitter phase ($y \rightarrow \rho_e/(4\Lambda)$).
3. The third possibility is characterised by $\delta < 0$ and $\Lambda < 0$ indicating $x < 0$ and $\rho_e > 0$. We did not find a reasonable solution in this case, although for $y > 0$ a $y \approx -x$ phase exists, but it cannot end in a low-energy accelerating phase with $x \rightarrow 0$ because all terms in Eq. (27) are positive in this limit. Hence, we do not pursue this solution any further.
4. Finally, the last case with $\delta > 0$ and $\Lambda > 0$ implies $x > 0$, and for $y < 0$ we




Figure 3: The numerical solutions from Eq. (31) for the scale factor $a(t)$, the Hubble rate $H(t)$ and the deceleration factor $q = -\ddot{a}/H^2$ for the case $\delta = 5 \cdot 10^6$, $\Lambda < 0$ described in Fig. 2.




Figure 4: The solution $y(x)$ from Eq. (27) and the functions $L_1(x)$ and $L_{12}(x)$ in the connection in Eqs. (30) for the case $\delta = -5 \cdot 10^6$, $\Lambda > 0$ and two values of the EOS $\omega = \frac{1}{3}, 0$.




Figure 5: The numerical solutions from Eq. (31) for the scale factor $a(t)$, the Hubble rate $H(t)$ and the deceleration factor $q = -\ddot{a}/H^2$ for the case $\delta = -5 \cdot 10^6$, $\Lambda > 0$ described in Fig. 4.

find a radiation/matter phase described by $y \approx -x$ for $x \rightarrow \infty$. Similar to the third case, $\rho_e > 0$ does not seem to allow a late-time de Sitter solution, however, the last term $\propto W^{-2}$ in Eq. (27) is positive giving rise to an accelerating solution with $x \rightarrow 0$. This cosmos is, however, quite exotic, because there is a contraction phase between the decelerating and the final de Sitter phase. For completeness the expansion behaviour of this interesting but unrealistic solution is shown in Fig. 6.

In the examples shown in this section we used relatively small values of $|\delta|$ for numerical reasons. According to its definition in Eq. (26), $\delta \sim (\Lambda/\rho_e)^{7/3}$ is related to the ratio of the large CC and the late-time energy density implying an enormous magnitude for δ . However, this does not correspond to unreasonable values for the parameter β in the action (20), instead a range of common values can be obtained depending on m and κ . Moreover, in this work we have normalised energy densities like r and κR by the constant Λ such that our results are independent of the numerical values of κ and Λ . For details on β and the relation between κ and Newton's constant we refer the reader to Ref. [33].

In summary, we found cosmological solutions for both signs of the large CC Λ , where the initial big bang singularity is replaced by a cosmic bounce with finite curvature and finite matter energy density. After the bounce these universes behave like being radiation or matter dominated, where the deceleration q is close to 1 or $\frac{1}{2}$, respectively. Eventually, when the matter component is sufficiently diluted the final de Sitter era begins, with an effective vacuum energy density $\sim \rho_e$ much smaller in magnitude than the large CC Λ . The latter property and the absence of an initial cosmological singularity are the main benefits of the CC filter model.




Figure 6: The solution $x(y)$ from Eq. (27), the scale factor $a(t)$ and the Hubble rate $H(t)$ from the numerical solution of Eq. (31) for the exotic case $\delta = 5000$, $\Lambda > 0$. Notice the contraction phase between the decelerating and accelerating eras.

5 Conclusions

The filter model for a large CC in the context of Palatini modified gravity [33] makes gravity insensitive to vacuum energy density contributions. In other words, the curvature is dominated by ordinary matter, whereas energy sources with the EOS of a CC do not contribute at leading order despite having a much larger energy density at late times.

In this work, we have explored the CC filter model in the context of the very early universe, where in classical theories one usually expects to find an initial big bang singularity at very high energies. However, in our setup we observed that the curvature singularity is replaced by a cosmic bounce with the matter energy density, the scale factor and the curvature remaining finite. In the case shown in Fig. 1 the boundedness can be seen already from the solution of the algebraic equation (27), whereas the other cases show the same feature after solving the differential equation (31). Moreover, our numerical results also confirm the analytical study of the late-time evolution in Ref. [33].

The absence of an initial space-time singularity could mean that our classical model for gravity does not require an ultraviolet completion from a theory of quantum gravity. However, in the light of the exotic form of the gravity action functional (20) it is unlikely that this setup is a fundamental theory. On the other hand, we may follow the argumentation in Ref. [54], where Palatini gravity models are proposed as effective continuum descriptions of loop quantum cosmology. In both approaches the regularisation of the initial singularity does not require the introduction of new degrees of freedom making them minimal solutions to this problem.

Independent of its theoretical origin, the CC filter model has the capability to handle two fundamental problems of gravity, the old CC problem and the avoidance of an initial singularity. Therefore, it is an attractive scenario for further investigations. In addition to the qualitative features discussed so far, it will be interesting to study more aspects of the CC filter model in the future, e.g. black hole or star interiors or anisotropies and inhomogeneities in cosmology.

Acknowledgements

This work has been supported in part by MEC and FEDER under project FPA2010-20807, by the Spanish Consolider-Ingenio 2010 program CPAN CSD2007-00042 and by DIUE/CUR Generalitat de Catalunya under project 2009SGR502.

References

- [1] S. Weinberg, “The cosmological constant problem,” *Rev. Mod. Phys.* **61** (1989) 1–23.
- [2] T. Padmanabhan, “Cosmological constant: The weight of the vacuum,” *Phys. Rept.* **380** (2003) 235–320, [arXiv:hep-th/0212290](#).
- [3] O. Bertolami, “The cosmological constant problem: a user’s guide,” *Int. J. Mod. Phys.* **D18** (2009) 2303–2310, [arXiv:0905.3110 \[gr-qc\]](#).
- [4] F. Bauer, J. Sola, and H. Stefancic, “Dynamically avoiding fine-tuning the cosmological constant: the ‘Relaxed Universe’,” *JCAP* **1012** (2010) 029, [arXiv:1006.3944 \[hep-th\]](#).
- [5] F. Bauer, “The Cosmological constant and the relaxed universe,” *J.Phys.Conf.Ser.* **259** (2010) 012083, [arXiv:1010.1106 \[hep-th\]](#).
- [6] S. M. Carroll, “The cosmological constant,” *Living Rev. Rel.* **4** (2001) 1, [arXiv:astro-ph/0004075](#).
- [7] P. J. E. Peebles and B. Ratra, “The cosmological constant and dark energy,” *Rev. Mod. Phys.* **75** (2003) 559–606, [arXiv:astro-ph/0207347](#).
- [8] S. Nojiri and S. D. Odintsov, “Introduction to modified gravity and gravitational alternative for dark energy,” *ECONF C0602061* (2006) 06, [arXiv:hep-th/0601213](#).
- [9] E. J. Copeland, M. Sami, and S. Tsujikawa, “Dynamics of dark energy,” *Int. J. Mod. Phys.* **D15** (2006) 1753–1936, [arXiv:hep-th/0603057](#).
- [10] E. Elizalde and D. Saez-Gomez, “F(R) cosmology in presence of a phantom fluid and its scalar-tensor counterpart:towards a unified precision model of the universe evolution,” *Phys. Rev.* **D80** (2009) 044030, [arXiv:0903.2732 \[hep-th\]](#).
- [11] R. Durrer, “What do we really know about Dark Energy?,” [arXiv:1103.5331 \[astro-ph.CO\]](#).
- [12] M. Li, X.-D. Li, S. Wang, and Y. Wang, “Dark Energy,” *Commun.Theor.Phys.* **56** (2011) 525–604, [arXiv:1103.5870 \[astro-ph.CO\]](#).
- [13] D. Saez-Gomez, “On Friedmann-Lemaître-Robertson-Walker cosmologies in non-standard gravity,” [arXiv:1104.0813 \[hep-th\]](#). Ph.D.Thesis (Advisor: Dr. Emilio Elizalde Rius, Dr. Sergei D. Odintsov).
- [14] F. Bauer, J. Sola, and H. Stefancic, “The Relaxed Universe: towards solving the cosmological constant problem dynamically from an effective action functional of gravity,” *Phys. Lett.* **B688** (2010) 269–272, [arXiv:0912.0677 \[hep-th\]](#).
- [15] J. Sola, “Cosmologies with a time dependent vacuum,” *J. Phys. Conf. Ser.* **283** (2011) 012033, [arXiv:1102.1815 \[astro-ph.CO\]](#).
- [16] F. Bauer, J. Sola, and H. Stefancic, “Relaxing a large cosmological constant in the astrophysical domain,” [arXiv:1105.1030 \[gr-qc\]](#). *to be published in Mod. Phys. Lett. A*.
- [17] H. Stefancic, “The solution of the cosmological constant problem from the inhomogeneous equation of state - a hint from modified gravity?,” *Phys. Lett.* **B670** (2009) 246–253, [arXiv:0807.3692 \[gr-qc\]](#).

- [18] F. Bauer, J. Sola, and H. Stefancic, “Relaxing a large cosmological constant,” *Phys. Lett.* **B678** (2009) 427–433, [arXiv:0902.2215 \[hep-th\]](#).
- [19] F. Bauer, “Perturbations in the relaxation mechanism for a large cosmological constant,” *Class. Quant. Grav.* **27** (2010) 055001, [arXiv:0909.2237 \[gr-qc\]](#).
- [20] A. Bonanno and M. Reuter, “Cosmology with self-adjusting vacuum energy density from a renormalization group fixed point,” *Phys. Lett.* **B527** (2002) 9–17, [arXiv:astro-ph/0106468](#).
- [21] S. Nobbenhuis, “Categorizing Different Approaches to the Cosmological Constant Problem,” *Found. Phys.* **36** (2006) 613–680, [arXiv:gr-qc/0411093](#).
- [22] S. M. Barr, S.-P. Ng, and R. J. Scherrer, “Classical cancellation of the cosmological constant reconsidered,” *Phys. Rev.* **D73** (2006) 063530, [arXiv:hep-ph/0601053](#).
- [23] F. K. Diakonos and E. N. Saridakis, “A Statistical Solution to the Cosmological Constant Problem in the Brane world,” *JCAP* **0902** (2009) 030, [arXiv:0708.3143 \[hep-th\]](#).
- [24] F. R. Klinkhamer and G. E. Volovik, “Self-tuning vacuum variable and cosmological constant,” *Phys. Rev.* **D77** (2008) 085015, [arXiv:0711.3170 \[gr-qc\]](#).
- [25] P. Batra, K. Hinterbichler, L. Hui, and D. N. Kabat, “Pseudo-redundant vacuum energy,” *Phys. Rev.* **D78** (2008) 043507, [arXiv:0801.4526 \[hep-th\]](#).
- [26] C. Charmousis, E. J. Copeland, A. Padilla, and P. M. Saffin, “General second order scalar-tensor theory, self tuning, and the Fab Four,” [arXiv:1106.2000 \[hep-th\]](#).
- [27] G. Dvali, S. Hofmann, and J. Khoury, “Degravitation of the cosmological constant and graviton width,” *Phys. Rev.* **D76** (2007) 084006, [arXiv:hep-th/0703027](#).
- [28] R. Erdem, “A way to get rid of cosmological constant and zero point energy problems of quantum fields through metric reversal symmetry,” *J. Phys.* **A41** (2008) 235401, [arXiv:0712.2989 \[hep-th\]](#).
- [29] S. P. Patil, “Degravitation, Inflation and the Cosmological Constant as an Afterglow,” *JCAP* **0901** (2009) 017, [arXiv:0801.2151 \[hep-th\]](#).
- [30] D. A. Demir, “Vacuum Energy as the Origin of the Gravitational Constant,” *Found. Phys.* **39** (2009) 1407–1425, [arXiv:0910.2730 \[hep-th\]](#).
- [31] D. A. Demir, “Stress-Energy Connection and Cosmological Constant Problem,” *Phys. Lett.* **B701** (2011) 496–502, [arXiv:1102.2276 \[hep-th\]](#).
- [32] C. P. Burgess and L. van Nierop, “Technically Natural Cosmological Constant From Supersymmetric 6D Brane Backreaction,” [arXiv:1108.0345 \[hep-th\]](#).
- [33] F. Bauer, “Filtering out the cosmological constant in the Palatini formalism of modified gravity,” *Gen. Rel. Grav.* **43** (2011) 1733–1757, [arXiv:1007.2546 \[gr-qc\]](#).
- [34] D. N. Vollick, “On the viability of the Palatini form of $1/R$ gravity,” *Class. Quant. Grav.* **21** (2004) 3813–3816, [arXiv:gr-qc/0312041](#).
- [35] T. P. Sotiriou, “The nearly Newtonian regime in Non-Linear Theories of Gravity,” *Gen. Rel. Grav.* **38** (2006) 1407–1417, [arXiv:gr-qc/0507027](#).
- [36] E. Barausse, T. P. Sotiriou, and J. C. Miller, “Curvature singularities, tidal forces and the viability of Palatini $f(R)$ gravity,” *Class. Quant. Grav.* **25** (2008) 105008, [arXiv:0712.1141 \[gr-qc\]](#).
- [37] T. P. Sotiriou and V. Faraoni, “ $f(R)$ Theories Of Gravity,” *Rev. Mod. Phys.* **82** (2010) 451–497, [arXiv:0805.1726 \[gr-qc\]](#).
- [38] G. J. Olmo, “Palatini Approach to Modified Gravity: $f(R)$ Theories and Beyond,” *Int. J. Mod. Phys.* **D20** (2011) 413–462, [arXiv:1101.3864 \[gr-qc\]](#).

- [39] G. Allemandi, A. Borowiec, and M. Francaviglia, “Accelerated cosmological models in Ricci squared gravity,” *Phys. Rev.* **D70** (2004) 103503, [arXiv:hep-th/0407090](#).
- [40] B. Li, J. D. Barrow, and D. F. Mota, “The Cosmology of Ricci-Tensor-Squared Gravity in the Palatini Variational Approach,” *Phys. Rev.* **D76** (2007) 104047, [arXiv:0707.2664 \[gr-qc\]](#).
- [41] B. Li, D. F. Mota, and D. J. Shaw, “Indistinguishable Macroscopic Behaviour of Palatini Gravities and General Relativity,” *Class. Quant. Grav.* **26** (2009) 055018, [arXiv:0801.0603 \[gr-qc\]](#).
- [42] B. Li, D. F. Mota, and D. J. Shaw, “Microscopic and Macroscopic Behaviors of Palatini Modified Gravity Theories,” *Phys. Rev.* **D78** (2008) 064018, [arXiv:0805.3428 \[gr-qc\]](#).
- [43] Q. Exirifard and M. M. Sheikh-Jabbari, “Lovelock Gravity at the Crossroads of Palatini and Metric Formulations,” *Phys. Lett.* **B661** (2008) 158–161, [arXiv:0705.1879 \[hep-th\]](#).
- [44] F. Bauer and D. A. Demir, “Inflation with Non-Minimal Coupling: Metric vs. Palatini Formulations,” *Phys. Lett.* **B665** (2008) 222–226, [arXiv:0803.2664 \[hep-ph\]](#).
- [45] M. Borunda, B. Janssen, and M. Bastero-Gil, “Palatini versus metric formulation in higher curvature gravity,” *JCAP* **0811** (2008) 008, [arXiv:0804.4440 \[hep-th\]](#).
- [46] S. Capozziello, M. De Laurentis, and V. Faraoni, “A bird’s eye view of $f(R)$ -gravity,” [arXiv:0909.4672 \[gr-qc\]](#).
- [47] A. De Felice and S. Tsujikawa, “ $f(R)$ theories,” *Living Rev. Rel.* **13** (2010) 3, [arXiv:1002.4928 \[gr-qc\]](#).
- [48] L. Amendola, K. Enqvist, and T. Koivisto, “Unifying Einstein and Palatini gravities,” *Phys. Rev.* **D83** (2011) 044016, [arXiv:1010.4776 \[gr-qc\]](#).
- [49] V. Vitagliano, T. P. Sotiriou, and S. Liberati, “The dynamics of metric-affine gravity,” *Annals Phys.* **326** (2011) 1259–1273, [arXiv:1008.0171 \[gr-qc\]](#).
- [50] T. Harko, T. S. Koivisto, and F. S. N. Lobo, “Palatini formulation of modified gravity with a nonminimal curvature-matter coupling,” *Mod. Phys. Lett.* **A26** (2011) 1467–1480, [arXiv:1007.4415 \[gr-qc\]](#).
- [51] F. Bauer and D. A. Demir, “Higgs-Palatini Inflation and Unitarity,” *Phys. Lett.* **B698** (2011) 425–429, [arXiv:1012.2900 \[hep-ph\]](#).
- [52] C. Barragan and G. J. Olmo, “Isotropic and Anisotropic Bouncing Cosmologies in Palatini Gravity,” *Phys. Rev.* **D82** (2010) 084015, [arXiv:1005.4136 \[gr-qc\]](#).
- [53] G. J. Olmo, “Palatini Actions and Quantum Gravity Phenomenology,” [arXiv:1101.2841 \[gr-qc\]](#).
- [54] G. J. Olmo and P. Singh, “Effective Action for Loop Quantum Cosmology a la Palatini,” *JCAP* **0901** (2009) 030, [arXiv:0806.2783 \[gr-qc\]](#).
- [55] R. P. Woodard, “Avoiding dark energy with $1/R$ modifications of gravity,” *Lect. Notes Phys.* **720** (2007) 403–433, [arXiv:astro-ph/0601672](#).
- [56] V. Vitagliano, T. P. Sotiriou, and S. Liberati, “The dynamics of generalized Palatini Theories of Gravity,” *Phys. Rev.* **D82** (2010) 084007, [arXiv:1007.3937 \[gr-qc\]](#).
- [57] G. J. Olmo, H. Sanchis-Alepuz, and S. Tripathi, “Dynamical Aspects of Generalized Palatini Theories of Gravity,” *Phys. Rev.* **D80** (2009) 024013, [arXiv:0907.2787 \[gr-qc\]](#).